

## **General Disclaimer**

### **One or more of the Following Statements may affect this Document**

- This document has been reproduced from the best copy furnished by the organizational source. It is being released in the interest of making available as much information as possible.
- This document may contain data, which exceeds the sheet parameters. It was furnished in this condition by the organizational source and is the best copy available.
- This document may contain tone-on-tone or color graphs, charts and/or pictures, which have been reproduced in black and white.
- This document is paginated as submitted by the original source.
- Portions of this document are not fully legible due to the historical nature of some of the material. However, it is the best reproduction available from the original submission.

NATIONAL AERONAUTICS AND SPACE ADMINISTRATION

**N84-11563**

Unclas  
G3/43 15221

## TABLE OF CONTENTS

	<u>Page</u>
LIST OF TABLES . . . . .	ii
LIST OF FIGURES. . . . .	iv
ABSTRACT . . . . .	1
INTRODUCTION . . . . .	2
EXPERIMENTAL . . . . .	5
RESULTS AND DISCUSSION . . . . .	6
CONCLUSION . . . . .	10
REFERENCES . . . . .	12

# LIST OF TABLES

	<u>Page</u>
Table 1	14
Concentration Changes of Free and Adsorbed Humic Acids with Time	
(a) Freshly Prepared Hydrous Iron Oxide. Adsorption Experiment at 35°C.	14
(b) Hydrous Iron Oxide in (a) Aged for Six Weeks. Adsorption Experiment at 35°C.	14
(c) Hydrous Iron Oxide of Seed I. Adsorption Experiment at 35°C.	15
(d) Freshly Prepared Hydrous Iron Oxide. Adsorption Experiment at 25°C.	15
Table 2	16
Adsorption of Humic Acids on Hydrous Iron Oxide in Fresh Water at 25°C. Expt. No. F1.	
Table 3	16
Adsorption of Humic Acids on Hydrous Iron Oxide in Fresh Water at 25°C. Expt. No. F2.	
Table 4	17
Adsorption of Humic Acids on Hydrous Iron Oxide in Fresh Water at 25°C. Expt. No. F3.	
Table 5	17
Adsorption of Humic Acids on Hydrous Iron Oxide in Fresh Water at 25°C. Expt. No. F4.	
Table 6	18
Adsorption of Humic Acids on Hydrous Iron Oxide in Fresh Water at 25°C. Expt. No. F5.	
Table 7	19
Adsorption of Humic Acids on Hydrous Iron Oxide in Seawater at 35°C. Expt. No. SW1.	
Table 8	19
Adsorption of Humic Acids on Hydrous Iron Oxide in Seawater at 35°C. Expt. No. SW2.	
Table 9	20
Adsorption of Humic Acids on Hydrous Iron Oxide in Seawater at 35°C. Expt. No. SW3.	
Table 10	20
Adsorption of Humic Acids on Hydrous Iron Oxide in Seawater at 35°C. Expt. No. SW4.	
Table 11	21
Adsorption of Humic Acids on Hydrous Iron Oxide in Seawater at 35°C. Expt. No. SW5.	
Table 12	21
Adsorption of Humic Acids on Hydrous Iron Oxide Seed I in Seawater at 35°C. Expt. No. SD1	
Table 13	22
Adsorption of Humic Acids on Hydrous Iron Oxide Seed III in Seawater at 25°C. Expt. No. SDIII-1.	
Table 14	22
Adsorption of Humic Acids on Hydrous Iron Oxide Seed III in Seawater at 25°C. Expt. No. SDIII-2.	

# LIST OF TABLES (con'd)

		<u>Page</u>
Table 15	Adsorption of Humic Acids on Hydrous Iron Oxide Seed V in Seawater at 35°C. Expt. No. SDV-1.	23
Table 16	Adsorption of Humic Acids on Hydrous Iron Oxide Seed V in Fresh Water at 35°C. Expt. No. SDV-2.	23
Table 17	Adsorption of Humic Acids on Hydrous Iron Oxide Seed V in Fresh Water at 35°C. Expt. No. SDV-3.	24
Table 18	Adsorption of Humic Acids on Hydrous Iron Oxide at 25°C and 35°C. Surface Concentration, $\Gamma$ , on Various Seeds With and Without Aging.	25
Table 19	Adsorption of Humic Acids on Hydrous Iron Oxide at 35°C. Surface Concentration, $\Gamma$ , on Various Seeds of Different Sizes.	25
Table 20	Adsorption Parameters of Humic Acids on Hydrous Iron Oxide.	26
Table 21	Adsorption of Cadmium on Hydrous Iron Oxide Seed V in Fresh Water at 35°C. Expt. No. Cd1.	27
Table 22	Adsorption of Cadmium of Hydrous Iron Oxide Seed V in the Presence of Humic Acid in Fresh Water at 35°C. Expt. No. Cd2.	27
Table 23	Adsorption of Cadmium on Hydrous Iron Oxide Seed V in Seawater at 35°C. Expt. No. Cd3.	28

# LIST OF FIGURES

		<u>Page</u>
Figure 1	Plot of Surface Concentration, $\Gamma$ Against Time. (o) Expt. A (◐) Expt. SD1 (∇) Expt. B1 (Δ) Expt. B2	30
Figure 2	Adsorption Isotherm in Fresh Water at 25°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids $C_f$ at Equilibrium. Expt. F5.	31
Figure 3	Adsorption Isotherms in Fresh Water at 25°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. (o) Expt. F1 (x) Expt. F2 (Δ) Expt. F3	32
Figure 4	Adsorption Isotherm in Fresh Water at 25°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. Expt. F4.	33
Figure 5	Langmuir Isotherm. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. F5.	34
Figure 6	Langmuir Isotherms. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . (o) Expt. F1 (x) Expt. F2 (Δ) Expt. F3.	35
Figure 7	Langmuir Isotherm. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. F4.	36
Figure 8	Adsorption Isotherms in Seawater at 35°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. (Δ) Expt. SW1 (o) Expt. SW2 (◐) Expt. SW3	37
Figure 9	Langmuir Isotherms. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . (Δ) Expt. SW1 (o) Expt. SW2 (◐) Expt. SW3	38

# LIST OF FIGURES (con'd)

	<u>Page</u>
Figure 10    Langmuir Isotherms. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . (o) Expt. SW2 ( $\square$ ) Expt. SW3	39
Figure 11    Adsorption Isotherms in Seawater at 35°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. Expt. SW4.	40
Figure 12    Langmuir Isotherm. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. SW4.	41
Figure 13    Adsorption Isotherm in Seawater at 25°C. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. Expt. SW5.	42
Figure 14    Langmuir Isotherm. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. SW5.	43
Figure 15    Adsorption Isotherms at 35°C for Hydrous Iron Oxide Seed V. Plot of Surface Concentration, $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. ( $\Delta$ ) Expt. SDV-1 ( $\square$ ) Expt. SDV-2 (o) Expt. SDV-3	44
Figure 16    Langmuir Isotherms for Seed V. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . ( $\Delta$ ) Expt. SDV-1 ( $\square$ ) Expt. SDV-2 (o) Expt. SDV-3	45
Figure 17    Adsorption Isotherms at 25°C for Hydrous Iron Oxide Seed III. Plot of Surface Concentration $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. Expt. SDIII.	46
Figure 18    Langmuir Isotherm for Seed III. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. SDIII.	47

# LIST OF FIGURES (con'd)

		<u>Page</u>
Figure 19	Adsorption Isotherm at 35°C for Hydrous Iron Oxide Seed I. Plot of Surface Concentration $\Gamma$ Against Free Humic Acids, $C_f$ at Equilibrium. Expt. SDI-2.	48
Figure 20	Langmuir Isotherm for Seed I. Plot of Surface Concentration, $\Gamma$ Against $\Gamma/C_f$ . Expt. SDI-2.	49
Figure 21	Adsorption Isotherm of Cadmium at 35°C. Plot of Surface Concentration, $\Gamma$ Against Free Cadmium, $C_f$ , at Equilibrium. (o) Expt. Cd1 ( $\Delta$ ) Expt. Cd2 ( $\square$ ) Expt. Cd3	50



## ABSTRACT

Studies concerning the interactions between suspended hydrous iron oxide and dissolved humic acids and trace metals are reported. As a major component of dissolved organic matters and its readiness for adsorption at the solid/water interface, humic acids may play a very important role in the organo-metallic geochemistry of suspended sediments and in determining the fate and distribution of trace metals, pesticides and anions in natural water systems.

Most of the solid phases in natural waters contain oxides and hydroxides. The most simple promising theory to describe the interactions of hydrous iron oxide interface is the surface complex formation model. In this model, the adsorptions of humic acids on hydrous iron oxide may be interpreted as complex formation of the organic bases (humic acid oxyanions,  $RO_x^{-z}$ ) with surface Fe ions. Measurements on adsorptions were made in both fresh water and seawater. Attempts have been made to fit our data to Langmuir adsorption isotherm. Adsorption equilibrium constants,  $K_1$ , were determined.

## INTRODUCTION

The presence of suspended particles in natural waters has a significant influence over the chemical composition of the whole system. The objective of our research project is to study the interactions between suspended particles and dissolved organic matters such as humic acids and trace metals.

Suspended solids are introduced into the natural bodies of water by natural forces, e.g. weathering, volcanic emanation, or by man-made activities, e.g. sewage discharge, dredging, continental drilling, ... Regardless of their sources and origins, suspended solids can be classified as inorganic oxides (quartz, ferric oxides,  $Al_2O_3$ ), carbonates, clays, and as organic detritus (organic colloidal matter of detrital origin), as well as living micro-organisms—algae and bacteria. The importance of suspended particles lies in their unique characters, specially small size, large specific surface area, and charge-carrying in the aqueous system. Solids in natural waters have electrically charged surfaces. Near neutral pH value, clay, most insoluble oxides, organic pollutants, bacteria and algae are characterized by a negative surface potential as indicated by negative values of the electrophoretic mobility.<sup>(1-4)</sup>

It is generally known that solutes tend to accumulate at the solid-solution interface. The concentration of chemical species, such as trace metals, hydrocarbon, phosphate and silicate in the bottom sediments or in the particulate solids of aqueous system is at least one or more orders of magnitude larger than that found in the solution part. The absorption characteristics of soluble chemicals, e.g. trace metals, pesticides, and inorganic anions onto various solids, e.g. oxides, carbonates, clays, soils, and organic detritus exactly reflect the importance of solid-solution interface and its role in controlling the chemical composition of the natural waters. All available information indicates that pH, the chemical composition, and crystal structure of the solid and the concentration of adsorbate govern significantly the

adsorption of metal ions at the solid-solution interface. A significant fraction of trace metals in water and sediment phase is often associated with organic matters such as humic substances (of the humic acid and fulvic acid type). Humic substances are highly oxidized and chemically and biologically stable polymeric compounds which have reported molecular weights in the range of several hundred to several million and carry carboxyl and hydroxyl functional groups. (5) As a major component of dissolved organic matters and its readiness for adsorption at the solid/water interface, humic acids may play a very important role in the organo-metallic geochemistry of suspended sediments and in determining the fate and distribution of trace metals, pesticides and anions in natural water systems.

Most of the solid phases in natural waters contain oxides or hydroxides. Various theories have been proposed to describe the absorption of ions at hydrous oxide interface. But the most simple and promising one is the surface complex formation model. (6-7) In this model the hydrolysis of oxide surface produces hydrous oxide surface groups as  $\text{>Si-OH}$ ,  $\text{>Mn-OH}$  and  $\text{>Fe-OH}$ . The surface hydroxyl groups have amphoteric properties. Protonation or deprotonation results in surface charge on the oxide surfaces, e.g.  $\text{>MeOH}_2^+ \rightleftharpoons \text{>MeOH} + \text{H}^+$   $\text{>MeOH} \rightleftharpoons \text{>MeO}^- + \text{H}^+$  where Me = metal (e.g. Fe, Mn, Si, Al) of solid oxides.

The pH of the solution is the parameter which determines the surface charge and potential of the oxides; and this solution property provides the mechanism through which the coulombic interactions between the ions and the surface may be controlled. The pH at which  $(\text{MeOH}_2^+) = (\text{MeO}^-)$  is called the pH of zero point of charge  $\text{pH}_{\text{ZPC}}$ . For example,  $\text{pH}_{\text{ZPC}}$  of Goethite ( $\alpha\text{-FeOOH}$ ) ranges from 7.5 to 8.3 in NaCl solution; (8) at oceanic pH of 8, the domination surface species is (FeOH).

Because of the pH-dependent charge characteristics, these oxides can exhibit either cation or anion exchange characteristics, dependent upon pH. But in

addition to their purely electrostatic effects on the ionic solutes, hydrous oxides show a strong tendency to interact chemically with anions as well as cations. These chemical interactions may be explained most satisfactorily in terms of complex formation. For example, the interactions with anions can be interpreted as a ligand exchange,  $\text{>Me-OH} + \text{HPO}_4^{2-} \longrightarrow \text{>Me-OPO}_3\text{H}^- + \text{OH}^-$ , and those with cations as a coordination with electron acceptors  $\text{>Me-OH} + \text{Cd}^{2+} \longrightarrow \text{>Me-OCd}^+ + \text{H}^+$ . In the natural water environment  $\gamma\text{-Al}_2\text{O}_3$  and goethite <sup>(9)</sup> are mostly responsible for the removal of Cd(II) and Zn(II).

During the past year we have made a number of measurements on the adsorption of humic acids on hydrous iron oxide mainly in fresh water and in seawater as well. The results of such adsorption experiments presented at the American Chemical Society's 34th Annual Southeastern Regional Meeting, Birmingham, Alabama, 1982, are shown in Figures 1-20 and Tables 1-20.

Although attempts have been made to prepare the same suspensions for adsorption studies from reagent solutions of the same concentrations, the results of such studies are very difficult to reproduce. The methods (such as rate of mixing, stirring, etc.) used to mix solutions for the preparation of these suspensions often influence the course of the subsequent precipitation, and so the availability of active sites on the solid surface as well as the size of the suspended particle. For these reasons, a seed suspension technique was adopted for some adsorption experiments so that we may be able to use the same kind of suspension for reproducible studies. The effects of factors such as ionic strength, temperature, pH and additives may be studied quantitatively. Also the influence of particle morphology may be investigated.

Undergraduate students were involved in this project to learn general techniques and the application of these techniques to the studies of adsorption in particular. As a result, the students have been familiar with (a) the use of atomic absorption spectrophotometry (AAS) for the analysis of trace metals;

(b) UV/visible spectrophotometer for the measurement of humic acids; (c) the performance of liquid scintillation counting techniques; (d) the separation of suspensions; and (e) the calculations and the principles of the interactions at the solid/liquid interface.

## EXPERIMENTAL

Adsorption experiments were designed to demonstrate under conditions similar to those in the natural water environments, the interactions of suspended particles with humic acids and the effect of adsorbed humic acids on trace metal uptake.

During some studies of adsorption in seawater, we used artificial rather than natural seawater to minimize biological effects and to provide a reproducible solution of known composition. The artificial seawater was prepared by the formula of Kester, et al.<sup>(10)</sup>

### Preparation of Solids (Adsorbents)

Amorphous iron oxide was prepared by slowly adding NaOH (2.5 M) to a vigorously stirred ferric nitrate solution until a pH of 8.0 was reached. NaOH to  $\text{Fe}(\text{NO}_3)_3$  was at an OH:FE ratio of three. The precipitate was aged before beginning adsorption experiments. In order to obtain fresh precipitate, the suspension from mixing NaOH and  $\text{Fe}(\text{NO}_3)_3$  was simply diluted with water (or artificial seawater) to give concentration in the desired range. 0.005 M  $\text{Fe}(\text{NO}_3)_3$  was used for the preparations of suspensions and was freshly prepared each time.

Since such experiments were very difficult to reproduce, a study was made of the adsorption of humic acids using a reproducible seed technique. Three preparations of seed suspensions of hydrous iron oxide were used. They were all prepared by adding 2.5 M NaOH solution to 0.005 M  $\text{Fe}(\text{NO}_3)_3$  solution,

washed several times and aged. Seed suspension I was prepared by slowly adding dilute NaOH solution to 0.005 M  $\text{Fe}(\text{NO}_3)_3$  with constant stirring. Seed suspension III was prepared by slowly (drop-wise) adding 2.5 M NaOH solution to 0.005 M  $\text{Fe}(\text{NO}_3)_3$  without much stirring. Seed suspension V was prepared by quickly adding 2.5 M NaOH solution to 0.005 M  $\text{Fe}(\text{NO}_3)_3$  with constant stirring. They were of mean size of about 5  $\mu\text{m}$ .

### Adsorption Analysis

The adsorption experiments were carried out in the suspensions of hydrous iron oxide of approximately 50 mg per liter. Humic acid was added to achieve the desired initial solute concentration. The mixture was shaken in stoppered bottles on an automatic shaker for 24 hours to equilibrate the adsorption system. The suspended particles were then isolated by filtration (pore size 0.45  $\mu\text{m}$ ). The concentrations of humic acids remaining in the aliquots were analyzed by a spectrophotometer, Spectronic 21 (Bausch & Lomb). The adsorbances of the resulting solutions were measured at 365 nm. PA-720 HIAC Particle Size Analyzer was used to measure the size distribution of particular matters.

### RESULTS AND DISCUSSION


The results have illustrated the effect of temperature, aging, and size and effective surface area on the adsorption behavior of humic acids on hydrous iron oxide. From Figure 1 and Table 1, it appears that 90% adsorption took place in the first 2 hours, equilibrium was reached after 20 hours of shaking and remained so to 48 hours. Criteria for attainment of equilibria in adsorption (or reactions) at hydrated solid surfaces are often difficult to establish. After the initially rapid adsorption, ferric oxide surfaces continue to adsorb humic acids very slowly. The slow adsorption (or reactions) may arise from the structure rearrangement of oxide surface to expose effectively a greater number

of FeOH sites at the interface.

The results as illustrated in Figure 1 show that adsorption of humic acids decreased rapidly with rising temperature. From the thermodynamic point of view, all adsorption processes are exothermic (i.e.,  $\Delta H$  must be negative), the extent of adsorption, therefore, decreases with increasing temperature.

The effects of aging hydrous iron oxide by allowing the suspensions to stand at room temperature for several weeks before adsorption experiments were studied. The results obtained showing a definite decrease in humic acids adsorption are presented in Figure 1 and Table 18. The aging effects appears very pronounced and cannot be ignored. The size relationship observation in Table 19 serves to demonstrate surface area control of humic acid adsorption, which were usually overlooked.


The adsorption of humic acids on hydrous iron oxide may be interpreted as complex formations of the organic bases (humic acid oxyanion,  $RO_x^{-z}$ ) with surface Fe ions. This is consistent with a mechanism involving ligand exchange of humic anion groups with  $H_2O$  and  $OH^-$  of surface  $Fe-OH_2^+$  and FeOH groups respectively. The adsorption of humic acids on hydrous iron oxide (and other suspended particles as well) is likely to modify substantially their surface properties in aqueous systems. The reaction between the oxyanions of humic acids and Fe of iron oxide surface is given



The bond strength between Fe and the O of oxyanion may be positively related to the logarithm of the complex formation constant  $K_1$  where

$$K_1 = [Fe-OR_{x-1}^{3-z}] / ([Fe^{3+}(aq)][RO_x^{-z}])$$

for a reaction.



However the complex formation constant,  $K_1$  is not yet known.

Adsorption is most often described in terms of isotherms which show the relationship between the quantity of solute adsorbed on suspended particles and bulk activity (concentration,  $C$ ) at a given temperature. Attempts have been made to fit our data to the Langmuir adsorption isotherm. Let the surface equilibrium for adsorption process be represented by the equation

$$1 - \theta + C_f \xrightleftharpoons{K_1} \theta$$

where  $C_f$  is the equilibrium concentration of adsorbate,  $\theta = \Gamma/\Gamma_m$ , the fractional coverage of surface sites;  $\Gamma$  is the adsorption density (adsorbed quantity) in mg per gram of suspended particles.  $\Gamma_m$  is the adsorption density for monolayer;  $K_1$  is the adsorption equilibrium constant related to the free energy of adsorption,  $\Delta G_{ads}$  by

$$K_1 = \exp (-\Delta G_{ads}^0/RT)$$

$$K_1 = \frac{\theta}{(1-\theta)C_f} = \frac{\Gamma/\Gamma_m}{(1-\Gamma/\Gamma_m)C_f}$$

$$\Gamma = \frac{\Gamma_m K_1 C_f}{1 + K_1 C_f}$$

The above Langmuir isotherm equation can be arranged into a linear form, thus

$$\Gamma = \Gamma_m - \left(\frac{1}{K_1}\right) \frac{\Gamma}{C_f}$$

The adsorption isotherms were determined by plotting (1)  $\Gamma$  versus  $C_f$  to show the change of adsorption (surface coverage) with free humic acids concentration. The results are given in Figures 2-4, 8, 11, 13, 15, 17, 19, and (2)  $\Gamma$  versus  $\Gamma/C_f$  gives a straight line with the slope of  $1/K_1$  and the intercept of  $\Gamma_m$ , as shown in Figures 5-7, 9, 10, 12, 14, 16, 18, 20. The values of adsorption capacity



$\Gamma_m$  (monolayer) and  $K_1$  (adsorption equilibrium constant) are summarized in Table 20. The results show good fits to the Langmuir isotherm.

From the experiment F1, F2 and F3 in which the suspensions were prepared under similar conditions, the parameters  $K_1 = \exp [-\Delta G_{\text{ads}}^0 / RT]$  appear to be closely the same. But the adsorption capacity,  $\Gamma_m$  (the number of available sites) are different from different suspensions. It may be that the reason for the different values of  $\Gamma_m$  is simply the size of suspended particles.

The most notable criticism of the Langmuir adsorption equation concerns the simplifying assumption that the free energy of adsorption  $\Delta G_{\text{ads}}$ , is independent of surface coverage ( $\theta$ ); solid surfaces tend to be non-uniform in nature. In our experiments, since the pH of the suspension during adsorption remained at about 8 ( $\pm 0.5$ ), the proportions of  $\text{Fe-OH}_2^+$ ,  $\text{Fe-OH}$  and  $\text{FeO}^-$  of the surface should remain about the same. In other words, the oxide surface remain in a similar environment, although the available adsorption sites decrease on adsorption. This may explain why  $K_1$  as a measure of the affinity of the surface of humic acids is more or less the same in experiments F1, F2, and F3 and so Langmuir equation is observed.

The results of the adsorption studies of  $\text{Cd}^{2+}$  on hydrous iron oxide are given in Figure 21 and Tables 21, 22, and 23. A nearly linear increase of adsorption either in fresh water or in seawater was observed over the range of free  $\text{Cd}^{2+}$  ion concentration from zero to 12  $\mu\text{g}$  per liter (when equilibrium was reached). The evidence indicates that  $\text{Cd}^{2+}$  ion was readily adsorbed by hydrous iron oxide at the value of pH equal 8. With the concentration of  $\text{Cd}^{2+}$  we used in this study, the adsorption capacity ( $\Gamma_m$ ) of hydrous iron oxide for  $\text{Cd}^{2+}$  ion was far from being reached since the limiting plateau of adsorption isotherms was out of sight. Results from the present study confirm the observation in literature<sup>(11)</sup> that dissolved trace metal concentrations in rivers are low and most likely non-problematic because more than 90% of the metal load of the river is associated

with particulate matter.

The effect of ionic strength on the adsorption was well demonstrated in Figure 21; the adsorption for  $\text{Cd}^{2+}$  was very much decreased in the medium of seawater. The ionic strength controls the charge density in the diffuse double layer. When the ionic strength is increased, at constant pH, the electrical double layer around the particle is compressed. The electric field ( $d\psi/dx$ ), at a given point, is increased, thus resulting in higher solvation energy contributions to the free energy of the adsorption process. The coulombic energy is also decreased, although the coulombic energy is minimized in hydrous iron oxide suspensions at pH equal 8. As a result, the adsorption of  $\text{Cd}^{2+}$  is reduced.

However, the solvation energy (as well as coulombic energy) decreases with increasing ionic dimensions. Thus, for humic acids because of large size of oxyanion, the increase of ionic strength becomes a favorable effect on the adsorption process as illustrated in Figure 15.

Since pH of Goethite ( $\alpha\text{-FeOH}$ ) ranges from 7.5 to 8.3 (Balistrieri and Murray, 1979)<sup>(8)</sup>, it is reasonable to assume that the specific interaction component (non-electrostatic) is the major contribution to the change of the Gibbs free energy,  $\Delta G_{\text{ads}}$ , during the adsorption on hydrous iron oxide. More adsorption experiments will be needed to illustrate possible mechanism. In addition to hydrous iron oxide, sediments collected from adjacent rivers may be used as an adsorbent in further study.

#### CONCLUSION

We present the results of adsorption experiments with humic acids and trace metal (Cd) using hydrous iron oxide as a model adsorbent for natural water systems. Although the concentrations of hydrous iron oxide are higher than normally would be encountered, the experimental conditions reflect the concentrations of humic acids (as dissolved organic carbon) and  $\text{Cd}^{2+}$  possibly found in

natural waters. The pH of all the reaction media was adjusted to 8.0, which closely represents the pH conditions of seawater.

Although the exact mechanism of surface bonding remains unknown, it is apparent that humic acids complexes are present at the surface. Since adsorbed humic acids possibly have functional groups directed toward solution, these ligands may serve as complexing agents at the surface. Humic acids coated on surface oxide may help control the distribution of trace metals in natural waters by binding through complexation with coated humic acids rather than reactions with simple oxide surface sites. In our study, the role of humic acids as a complexing ligand in the uptake of  $\text{Cd}^{2+}$  by hydrous iron oxide still remains largely inconclusive by the data in fresh water (Tables 21 and 22, and Figure 21). Further experiments are planned to study the adsorption of  $\text{Cd}^{2+}$  in seawater in the presence and absence of humic acids. Since in seawater, the suspended particles of hydrous iron oxide are better coated with humic acids, and so the effect of adsorbed humic acids on the uptake of  $\text{Cd}^{2+}$  may be better illustrated.

The adsorption data of humic acids and  $\text{Cd}^{2+}$  ion at pH equal 8.0 indicate that there is appreciable adsorption in the vicinity of PZC<sup>(8)</sup>. This behavior can be attributed only to the specific adsorption of humic acid and/or  $\text{Cd}^{2+}$  on hydrous iron oxide.

Since cadmium is a highly toxic pollutant, and often a component of industrial waste, it is useful to know whether this transition metal complex with humic acids is adsorbed. On the other hand, dissolved humic acids in solution may compete with the oxide surface for coordination of  $\text{Cd}^{2+}$  ions;  $\text{Cd}^{2+}$  adsorption may be decreased by complexing with the humic acids remaining in solution. This clearly points out the need for more research in order to solve problems of great significance.

## REFERENCES

1. S. D. Faust and M. C. Magner, Water and Sewage Works, 111, (1964).
2. G. Tchobanoglous and R. Eliassen, Jour. Sanitary Eng. Div., Proceedings of Amer. Soc. Civil Eng., SA2, 243, (1970).
3. R. A. Neihof and G. I. Loeb, Amer. Soc. Limnol. and Oceanogr., 17, 7, (1972).
4. R. A. Baker, S. J. Campbell, and F. R. Anspach, Jour. Amer. Water Works Assoc., 57, 363, (1965).
5. M. A. Rashid and L. H. King. Geochim. Cosmochim. ACTA, 34, 193-201 (1970).
6. W. Stumm and J. J. Morgan, Aquatic Chemistry, 583 pp, Wiley, New York (1970).
7. P. W. Schindler, Thalassia Jugosl., 11, 101-111, (1975).
8. L. Balistrieri and J. W. Murray, Surface of Goethite ( $\alpha\text{FeOOH}$ ) in Seawater. In Chemical Modeling in Aqueous Systems (ed., E. A. Henne), 914 pp. Amer. Chem. Soc., No. 93.
9. E. A. Forbes and J. P. Quirk, Jour. Colloid Interface Sci., 49(3), 403 (1974).
10. D. R. Kester, I. W. Duedall, D. N. Connors and R. M. Pytkowicz, Limnol. Oceanogr. 12, 176-179 (1967).
11. B. J. Presley, J. H. Trefry and R. F. Shokes, Water, Air, and Soil Pollution 13, 481-494 (1980).

## TABLES

TABLE 1: CONCENTRATION CHANGES OF FREE AND ADSORBED HUMIC ACIDS WITH TIME

(a): Freshly Prepared Hydrous Iron Oxide. Adsorption Experiment at 35°C.

	1	2	3	4	5	6	7	8
Initial Concentration of HA mg/l	8.67	8.67	8.67	8.67	8.67	8.67	8.67	8.67
Time Elapsed (Hour)	0.5	2.0	5.0	8.3	18.5	24	31	42.5
Free [HA] in Solution mg/l	6.22	4.60	2.77	1.22	1.02	0.93	0.87	0.93
[HA] Adsorbed on Iron Oxide mg/g	50.1	83.2	120.6	152.2	156.2	158.2	159.4	158.2

N.B. The concentration of suspension was 49.36 mg/l

(b): Hydrous Iron Oxide in (a) Aged for Six Weeks.  
Adsorption Experiment at 35°C.

	1	2	3	4	5	6	7
Initial Concentration of HA mg/l	8.67	8.67	8.67	8.67	8.67	8.67	8.67
Time Elapsed (Hour)	0.5	2.0	6.5	18.7	24.1	30.0	48.0
Free [HA] in Solution mg/l	4.25	3.07	2.17	1.75	1.47	1.48	1.25
[HA] Adsorbed on Iron Oxide mg/g	90.3	114.4	132.8	141.4	147.1	146.9	151.6

N.B. The concentration of suspension was 49.36 mg/l.

ORIGINAL PAGE  
OF POOR QUALITY

(c): Hydrous Iron Oxide of Seed I. Absorption Experiment at 35°C.

	1	2	3	4	5	6
Initial Concentration of HA mg/l	8.67	8.67	8.67	8.67	8.67	8.67
Time Elapsed (Hour)	0.5	1.5	4.0	8.5	20.0	26.0
Free [HA] in Solution mg/l	8.28	8.17	7.83	7.42	6.57	6.55
[HA] Adsorbed on Iron Oxide mg/g	8.25	10.8	17.8	26.7	44.6	44.8

N.B. The concentration of suspension was 49.36 mg/l

(d): Freshly Prepared Hydrous Iron Oxide.  
Adsorption Experiment at 25°C.

	1	2	3	4	5	6	7	8
Initial Concentration of HA mg/l	13.16	13.16	13.16	13.16	13.16	13.16	13.16	13.16
Time Elapsed (Hour)	0.5	1.5	4.5	9.5	19.0	25.0	31.5	42.5
Free [HA] in Solution mg/l	6.41	5.51	3.14	2.19	1.67	1.58	1.48	1.49
[HA] Adsorbed on Iron Oxide mg/g	138.6	157.0	205.7	225.2	235.9	237.7	239.8	239.6

N.B. The concentration of all above suspensions were 49.36 mg/l

ORIGINAL PAGE IS  
OF POOR QUALITY

ORIGINAL PAGE NO.  
OF POOR QUALITY

TABLE 2: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Fresh Water at 25°C - Expt. No. F1.

Reaction Flask No.	1	2	3	4	5	6
Free Humic Acid at Equilibrium $C_f$ mg/l	1.47	2.37	3.25	4.81	6.47	8.39
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	150.0	176.6	203.4	216.2	226.8	231.8
$\Gamma/C_f$	102.1	74.5	62.6	45.0	35.1	27.6

\*Freshly prepared.

TABLE 3: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Fresh Water at 25°C - Expt. No. F2.

Reaction Flask No.	1	2	3	4	5	6	7	8	9	10
Free Humic Acid at Equilibrium/ $C_f$ mg/l	1.01	1.67	1.77	2.90	4.17	4.20	5.88	6.07	7.45	9.17
Humic Acids Ad- sorbed on Iron Oxide $\Gamma$ mg/g	114.4	145.9	143.9	165.7	184.6	183.9	194.2	190.3	206.6	215.7
$\Gamma/C_f$	113.3	87.4	81.3	57.1	44.3	43.8	33.0	31.4	27.7	23.5

\*Freshly prepared.



ORIGINAL PAGE OF  
OF POOR QUALITY

TABLE 4: Adsorption of Humic Acids on Hydron Iron Oxide\*  
in Fresh Water at 25°C - Expt. No. F3.

Reaction Flask No.	1	2	3	4	5	6	7	8	9	10
Free Humic Acids at Equilibrium $C_f$ mg/l	1.30	1.35	2.23	3.22	3.35	5.28	5.38	6.75	6.82	8.3
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	108.5	107.5	134.5	159.2	156.5	161.8	159.7	176.3	174.9	187
$\Gamma/C_f$	83.5	79.6	60.3	49.4	46.7	30.6	29.7	26.1	25.6	22.

\*Freshly prepared.

TABLE 5: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Fresh Water at 25°C - Expt. No. F4.

Reaction Flask No.	1	2	3	4	5	6	7	8
Free Humic Acid at Equilibrium $C_f$ mg/l	0.52	1.13	1.90	3.17	4.38	5.58	7.77	9.60
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	71.0	120.0	141.2	160.2	180.2	200.4	200.0	206.8
$\Gamma/C_f$	137.0	99.1	74.3	50.5	41.2	35.9	25.7	21.6

\*Freshly prepared.

TABLE 6: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Fresh Water at 25°C - Expt. No. F5.

Reaction Flask No.	1	2	3	4	5	6	7	8	9	10	11
Free Humic Acids at Equilibrium, $C_f$ mg/l	0.69	0.69	1.30	1.30	2.08	3.92	4.05	5.42	5.42	6.17	8.19
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	120.9	120.9	153.5	153.5	182.5	189.7	187.0	203.7	203.7	188.2	191.3
$\Gamma/C_f$	175.3	175.3	118.1	118.8	87.7	48.4	46.2	37.6	37.6	30.5	23.4

\*Freshly prepared.

ORIGINAL PAGE IS  
OF POOR QUALITY

TABLE 7: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Sea Water at 35°C - Expt. No. SW1.

Reaction Flask No.	1	2	3	4	5	6
Free Humic Acids at Equilibrium, $C_f$ mg/L	0.12	0.32	0.73	1.69	3.56	5.95
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	58.1	94.4	126.0	146.5	168.1	178.9
$\Gamma/C_f$	484.5	295.0	172.6	86.7	47.2	30.1

\*Freshly prepared.

TABLE 8: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Sea Water at 35°C - Expt. No. SW2.

Reaction Flask No.	1	2	3	4	5	6
Free Humic Acid at Equilibrium $C_f$ mg/L	0.34	0.50	0.91	1.94	3.25	4.94
Humic Acid, Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	73.7	70.5	102.3	121.6	134.7	140.2
$\Gamma/C_f$	216.9	141.0	112.4	62.7	41.5	28.4

\*Freshly prepared.

TABLE 9: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Seawater at 35°C - Expt. No. SW3.

Reaction Flask No.	1	2	3	4	5
Free Humic Acid at Equilibrium $C_f$ mg/L	0.83	1.76	3.33	5.06	6.94
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	63.8	85.1	93.4	98.1	99.7
$\Gamma/C_f$	76.9	48.3	28.0	19.4	14.4

\*Freshly prepared.

TABLE 10: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Seawater at 35°C - Expt. No. SW4.

Reaction Flask No.	1	2	3	4	5
Free Humic Acid at Equilibrium $C_f$ mg/L	0.14	0.33	2.24	4.84	6.64
Humic Acid Ad- sorbed on Iron Oxide $\Gamma$ mg/g	37.7	73.9	115.5	142.2	145.2
$\Gamma/C_f$	269.3	224.1	51.6	29.4	21.9

TABLE 11: Adsorption of Humic Acids on Hydrous Iron Oxide\*  
in Seawater at 35°C - Expt. No. SW5.

Reaction Flask No.	1	2	3	4
Free Humic Acid at Equilibrium $C_f$ mg/l	0.50	1.35	2.89	4.28
Humic Acids Ad- sorbed on Iron Oxide $\Gamma$ mg/g	70.5	93.4	102.3	113.9
$\Gamma/C_f$	140.0	69.2	35.4	26.6

\*Freshly prepared.

TABLE 12: Adsorption of Humic Acids on Hydrous Iron Oxide  
Seed I in Seawater at 35°C. - Expt. No. SDI.

Reaction Flask No.	1	2	3	4
Free Humic Acid at Equilibrium $C_f$ mg/l	1.07	1.65	3.33	5.32
Humic Acids Ad- sorbed on Iron Oxide $\Gamma$ mg/g	34.0	41.3	46.6	46.4
$\Gamma/C_f$	31.8	25.0	14.0	8.7

TABLE 13: Adsorption of Humic Acids on Hydrous Iron Oxide Seed III in Seawater at 25°C - Expt. No. SDIII-1.

Reaction Flask No.	1	2	3	4	5	6	7
Free Humic Acid at Equilibrium $C_f$ mg/l	0.28	0.40	0.79	0.87	1.28	1.54	2.45
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	58.1	98.1	132.0	172.5	205.7	241.9	264.1
$\Gamma/C_f$	207.3	245.3	167.0	198.2	160.7	157.1	107.8

TABLE 14: Adsorption of Humic Acids on Hydrous Iron Oxide Seed III in Seawater at 25°C - Expt. No. SDIII-2.

Reaction Flask No.	1	2	3	4	5	6	7
Free Humic Acid at Equilibrium $C_f$ mg/l	0.30	0.47	0.68	0.92	1.39	1.85	2.34
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	57.6	96.6	134.3	171.4	203.3	235.2	266.5
$\Gamma/C_f$	182.1	205.6	197.5	186.3	146.3	127.2	113.9

TABLE 15: Adsorption of Humic Acids on Hydrous Iron Oxide  
Seed V in Seawater at 35°C - Expt. No. SDV-1.

Reaction Flask No.	1	2	3	4	5	6
Free Humic Acid at Equilibrium $C_f$ mg/l	0.27	0.38	0.50	0.70	0.88	1.02
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	88.3	123.2	157.9	190.9	224.1	258.1
$\Gamma/C_f$	327.2	324.3	315.8	272.8	254.7	253.0

Aged 24 hours.

TABLE 16: Adsorption of Humic Acids on Hydrous Iron Oxide  
Seed V\* in Fresh Water at 35°C - Expt. No. SDV-2.

Reaction Flask No.	1	2	3	4	5	6	7	8
Free Humic Acid Equilibrium $C_f$ mg/l	0.46	0.83	1.25	1.77	2.13	2.63	3.65	4.54
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	47.5	77.8	107.3	134.1	164.1	191.3	208.8	228.4
$\Gamma/C_f$	103.2	93.8	85.8	75.8	77.1	72.7	57.2	50.3

\*Aged 24 hours.

TABLE 17: Adsorption of Humic Acids on Hydrous Iron Oxide  
Seed V\* in Fresh Water at 35°C - Expt. No. SDV-3.

Reaction Flask No.	1	2	3	4	5
Free Humic Acids at Equilibrium $C_f$ mg/l	0.69	1.29	1.21	2.30	3.44
Humic Acids Ad- sorbed on Iron Oxide, $\Gamma$ mg/g	47.7	76.6	78.2	117.0	134.0
$\Gamma/C_f$	69.2	59.4	64.6	50.9	39.0

\*Aged for 8 weeks.



TABLE 18: Adsorption of Humic Acids on Hydrous Iron Oxide at 25°C and 35°C. Surface Concentrations  $\Gamma$ , on Various Seeds With and Without Aging.

	a	b	c	d
Hydrous Iron Oxide	Freshly prepared	From a. Aged 6 wks.	Seed I. Aged 5 wks.	Freshly prepared.
Temperature of Bath	35°C	35°C	35°C	25°C
Co, Initial Concentration of HA mg/l	8.67	8.67	8.67	13.16
$\Gamma$ [HA] Adsorbed on Iron Oxide at Equilibrium mg/g	158.4 mg/g	146.9	44.6	236.0

N.B. All concentrations of suspensions = 49.36 mg/l

TABLE 19: Adsorption of Humic Acids on Hydrous Iron Oxide at 35°C. Surface Concentrations  $\Gamma$ , on Various Seeds of Different Sizes.

	a	b
Hydrous Iron Oxide	Freshly prepared.	Freshly prepared
Temperature of Bath	35°C	35°C
Co, Initial Concentrations of HA mg/l	8.67	8.67
Sizes* of Particles (length in $\mu\text{m}$ )	4.2	4.7
$\Gamma$ [HA] Adsorbed on Iron Oxide at Equilibrium mg/g	7.75 158.37 mg/g	6.72 137.32 mg/g

N.B. All concentrations of suspensions = 49.36 mg/l.

\*PA-720 HIAC Particle Size Analyser was used.

TABLE 20: Adsorption Parameters of Humic Acids on Hydrous Iron Oxide.

Expt. No.	Temp. °C	Suspension Conc mg/	$\Gamma_m$	$K_1$	Correlation Co- efficient r	Avg. Size $\mu m$	Remarks
F1	25	48.8 $\pm$ 0.3	266	0.88	.989	-	Fresh Water
F2	"	"	231	0.95	.986	-	"
F3	"	"	206	0.85	.967	-	"
F4	"	"	231	0.86	.977	-	"
F5	"	"	214	1.94	.967	-	"
SW1	35	48.8 $\pm$ 0.3	177	3.88	.986	2.7	Sea Water
SW2	"	"	147	2.55	.929	6.2	"
SW3*	"	"	109	1.76	.991	4.9	"
SW4	"	"	149	2.61	.976	3.1	"
SW5	"	"	119	2.89	.981	-	"
SDI	35	55.0 $\pm$ 0.4	53	1.87	.945	-	Sea Water
SDIII-1	25	46.3 $\pm$ 0.3	435	0.64	.847	-	"
SDIII-2	"	"	435	0.64	.847	-	"
SDV-1	35	52.8 $\pm$ 0.3	614	0.67	.950	5.5	Sea Water
SDV-2	"	"	418	0.28	.967	"	Fresh Water
SDV-3*	"	"	248	0.36	.963	"	"

$K_1$  is adsorption equilibrium constant;  $\Gamma_m$  is monolayer adsorption, mg/g.

\*Seeds were aged for more than 6 weeks.

N.B. (1) Initial concentration of humic acids in suspensions were less than 15 mg/l to avoid coagulation when seawater used as medium.

(2) PA-720 HIAC Particle Size Analyser was used to estimate the average size of particles.

TABLE 21: Adsorption of Cadmium on Hydrous Iron Oxide  
Seed V in Fresh Water at 35°C - Expt. No. Cd1

	1	2	3	4	5	6
Free Cadmium at Equilibrium, $C_f$ , $\mu\text{g/l}$	0	0	.01	.04	.03	.07
Cadmium Adsorbed on Iron Oxide, $\Gamma$ $\mu\text{g/g}$	0	62.0	123.4	204.6	285.9	365.4

TABLE 22: Adsorption of Cadmium on Hydrous Iron Oxide  
Seed V in the Presence of Humic Acid in  
Fresh Water at 35°C - Expt. No. Cd2

	1	2	3	4	5	6
Free Cadmium at Equilibrium, $C_f$ , $\mu\text{g/l}$	0	.06	.14	.22	.37	.40
Cadmium Adsorbed on Iron Oxide, $\Gamma$ $\mu\text{g/g}$	0	60.8	120.7	200.8	278.8	358.5

TABLE 23: Adsorption of Cadmium on Hydrous Iron Oxide  
Seed V in Seawater at 35°C - Expt. No. Cd3

	1	2	3	4	5	6
Free Cadmium at Equilibrium $C_f$ , $\mu\text{g/g}$	0	2.15	3.93	6.22	9.33	11.64
Cadmium Adsorbed on Iron Oxide, $\Gamma$ , $\mu\text{g/g}$	0	17.4	42.1	76.3	92.9	125.3

## FIGURES




Fig. 1: Plot of Surface Concentration,  $\Gamma$  Against Time.  
( $\circ$ ) Expt. A; ( $\Delta$ ) Expt. SD-1; ( $\square$ ) Expt. B1; ( $\nabla$ ) Expt. B2




Fig. 2: Adsorption Isotherm in Fresh Water at 25°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. F5.




Fig. 3: Adsorption Isotherms in Fresh Water at 25°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. (○) Expt. F1; (×) Expt. F2; (Δ) Expt. F3.






Fig. 4: Adsorption Isotherm in Fresh Water at 25°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. F4




Fig. 5: Langmuir Isotherm. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
Expt. F5




Fig. 6: Langmuir Isotherms. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
 (o) Expt. F1; (x) Expt. F2; (Δ) Expt. F3




Fig. 7: Langmuir Isotherm. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
Expt. F4




Fig. 8: Adsorption Isotherms in Seawater at 35°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. (Δ) Expt. SW1; (○) Expt. SW2; (◻) Expt. SW3




Fig. 9: Langmuir Isotherms. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
 ( $\Delta$ ) Expt. SW1; ( $\circ$ ) Expt. SW2; ( $\square$ ) Expt. SW3




Fig. 10: Langmuir isotherms. Plot of surface concentration  $\Gamma$  against  $\Gamma/C_f$ .  
 (o) Expt. SW2; (□) Expt. SW3

ORIGINAL PAGE IS  
 OF POOR QUALITY




Fig. 11: Adsorption Isotherms in Seawater at 35°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. SW4






Fig. 12: Langmuir Isotherm. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
Expt. SW4




Fig. 13: Adsorption Isotherm in Seawater at 25°C. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. SW5




Fig. 14: Langmuir Isotherm. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
Expt. SW5




Fig. 15: Adsorption Isotherms at 35°C for Hydrous Iron Oxide Seed V. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium.  
 ( $\Delta$ ) Expt. SDV-1; ( $\square$ ) Expt. SDV-2; ( $\circ$ ) Expt. SDV-3




Fig. 16: Langmuir Isotherms for Seed V. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ .  
 ( $\Delta$ ) Expt. SDV-1; ( $\square$ ) Expt. SDV-2; ( $\circ$ ) Expt. SDV-3




Fig. 17: Adsorption Isotherms at 25°C for Hydrous Iron Oxide Seed III. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. SD III




Fig. 18: Langmuir Isotherm for Seed III. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ : Expt. SD III




Fig. 19: Adsorption Isotherm at  $35^{\circ}\text{C}$  for Hydrous Iron Oxide Seed I. Plot of Surface Concentration,  $\Gamma$  Against Free Humic Acids,  $C_f$  at Equilibrium. Expt. SDI-2






Fig. 20: Langmuir Isotherm for Seed I. Plot of Surface Concentration,  $\Gamma$  Against  $\Gamma/C_f$ . Expt. SDI-2




Figure 21: Adsorption Isotherm of Cadmium at 35°C. Plot of Surface Concentration,  $\Gamma$ , Against Free Cadmium,  $C_f$ , at Equilibrium.  
(o) Expt. Cd1; ( $\Delta$ ) Expt. Cd2; ( $\square$ ) Expt. Cd3